VOL. 494 JOURNAL OF CHROMATOGRAPHY SEPTEMBER 29, 1989

(Biomedical Applications, Vol. 86)

CONTENTS

(Abstracts/Contents Lists published in Analytical Abstracts, ASCA, Biochemical Abstracts, Biological Abstracts, Chemical Abstracts, Chemical Titles, Chromatography Abstracts, Clinical Chemistry Lookout, Current Contents/Physical, Chemical & Earth Sciences, Current Contents/Life Sciences, Deep-Sea Research/Part B: Oceanographic Literature Review, Excerpta Medica, Index Medicus, Mass Spectrometry Bulletin, PASCAL-CNRS, Pharmaceutical Abstracts, Referativnyi Zhurnal, Science Citation Index and Trends in Biotechnology)

Review. Potential of wide bore open tubular columns in gas chromatographic analysis of drugs
by A.C. Mehta, (Leeds, U.K.) (Received April 10th, 1989) 1

Simultaneous determination of the primary prostanoids prostaglandin E_2, prostaglandin $F_{2\alpha}$ and 6-oxoprostaglandin $F_{1\alpha}$ by immunoaffinity chromatography in combination with negative ion chemical ionization gas chromatography–tandem mass spectrometry
by G. Mackert (Heidelberg, F.R.G.), M. Reinke (Erlangen, F.R.G.) and H. Schweer and H.W. Seyberth (Heidelberg, F.R.G.) (Received April 17th, 1989) 13

Long-chain fatty acids and alcohols from gerbil meibomian lipids
by D.J. Harvey (Oxford, U.K.) (Received May 25th, 1989) 23

Improved determination of very-long-chain fatty acids in plasma and cultured skin fibroblasts: applications to the diagnosis of peroxisomal disorders
by W. Onkenhout, P.F.H. van der Poel and M.P.M. van den Heuvel (Leiden, The Netherlands) (Received May 4th, 1989) 31

Separation and quantitation of free cholesterol and cholesteryl esters in a macrophage cell line by high-performance liquid chromatography
by R. Vercaemst, A. Union and M. Rosseneu (Bruges, Belgium) (Received May 10th, 1989) 43

Catecholamine measurements in urine by high-performance liquid chromatography with amperometric detection — Comparison with an autoanalyser fluorescence method
by P. Hjemdahl, P.T. Larsson, T. Bradley, T. Åkerstedt, I. Anderzén, K. Sigurdsson, M. Gillberg and U. Lundberg (Stockholm, Sweden) (Received May 10th, 1989) 53

Measurement of methionine enkephalin and leucine enkephalin in rat brian regions by high-performance liquid chromatography with coulometric electrochemical detection
by C. Kim, R. Cheng and S.R. George (Toronto, Canada) (Received May 18th, 1989) 67

Determination of quinonoid dihydrobiopterin by high-performance liquid chromatography and electrochemical detection
by S. Heales and K. Hyland (London, U.K.) (Received May 27th, 1989) 77

Simple apparatus for capillary zone electrophoresis and its application to protein analysis
by V. Rohlíček and Z. Deyl (Prague, Czechoslovakia) (Received May 10th, 1989) .. 87

(Continued overleaf)

Contents (continued)

Capillary gas chromatographic–mass spectrometric determination of stable isotopically labelled and unlabelled theophylline in serum and urine and of 1,3-dimethyluric acid in urine
by Y. Kasuya, T. Furuta and H. Shimota (Tokyo, Japan) (Received May 23rd, 1989) 101

Sulfonium salts as derivatizing agents. 2. Determination of theophylline in plasma by automated gas chromatography
by B.L. Lee, P. Jacob, III and N.L. Benowitz (San Francisco, CA, U.S.A.) (Received May 30th, 1989) 109

Determination of the partial benzodiazepine receptor agonist Ro 16-6028 in plasma by capillary gas chromatography with nitrogen-selective detection after conversion into the ethyl ester derivative
by U. Timm, G. Fischer, M. Zell and R. Zumbrunnen (Basle, Switzerland) (Received April 7th, 1989) 119

Capillary gas chromatography of trihexyphenidyl, procyclidine and cycrimine in biological fluids
by J.A. Owen, M. Sribney, J.S. Lawson, N. Delva and F.J.J. Letemendia (Kingston, Canada) (Received May 5th, 1989) 135

Comparison between two methods for the determination of the total and free (*R*)- and (*S*)-disopyramide in plasma using an α_1-acid glycoprotein column
by M. Enquist and J. Hermansson (Stockholm, Sweden) (Received April 20th, 1989 143

Determination of (*R*)- and (*S*)-propranolol in plasma by high-performance liquid chromatography using N-benzoxycarbonylglycyl-L-proline as chiral selector in the mobile phase
by A. Karlsson and C. Pettersson (Uppsala, Sweden) and S. Björkman (Malmö, Sweden) (Received April 26th, 1989) 157

High-performance liquid chromatographic analysis of the enantiomers of flurbiprofen and its metabolites in plasma and urine
by M.P. Knadler and S.D. Hall (Indianapolis, IN, U.S.A.) (Received May 10th, 1989) 173

Quantification of L-3-(3-hydroxy-4-pivaloyloxyphenyl)alanine (NB-355) by high-performance liquid chromatography using *o*-phthalaldehyde/N-acetyl-L-cysteine derivatization
by A. Hisaka, S. Kasamatsu, N. Takenaga and M. Ohtawa (Tokyo, Japan) (Received May 26th, 1989) 183

High-performance liquid chromatographic method for the quantification of several diamidine compounds with potential chemotherapeutic value
by B.J. Berger, J.E. Hall and R.R. Tidwell (Chapel Hill, NC, U.S.A.) (Received May 3rd, 1989) 191

Comparison of high-performance liquid chromatography and the Abbott fluorescent polarization radioimmunoassay in the measurement of methotrexate
by W. Cosolo, O.H. Drummer and N. Christophidis (Heidelberg, Australia) (Received April 25th, 1989) 201

Determination of manidipine and its pyridine metabolite in human serum by high-performance liquid chromatography with ultraviolet detection and column switching
by T. Miyabayashi, K. Yamashita, I. Aoki, M. Motohashi, T. Yashiki and K. Yatani (Osaka, Japan) (Received April 20th, 1989) 209

Simultaneous determination of pyrimethamine and mefloquine in human plasma by high-performance liquid chromatography with ultraviolet detection
by A. Guenzi, G. Cappelletti, A. Scala and M. Zanetti (Milan, Italy) (Received May 3rd, 1989) 219

Analytical studies on β-lactam antibiotics. III. Automated high-performance liquid chromatographic method for the determination of the orally active antibiotic ceftibuten in human plasma and urine
by A. Matsuura, T. Nagayama and T. Kitagawa (Osaka, Japan) (Received May 17th, 1989) 231

Simultaneous determination of flurazepam and its metabolites in human plasma by high-performance liquid chromatography
by K. Selinger, D. Lessard and H.M. Hill (Senneville, Canada) (Received May 10th, 1989) 247

Determination of an oxadiazole-substituted 1,4-benzodiazepine in plasma by high-performance liquid chromatography with ultraviolet detection and by a radioreceptor assay
by L. Nordholm and H. Mengel (Soeborg, Denmark) (Received May 12th, 1989) ... 257

Sensitive analysis of the mycotoxin zearalenone and its metabolites in biological fluids by high-performance liquid chromatography
by D.B. Prelusky, R.M. Warner and H.L. Tenholm (Ottawa, Canada) (Received May 27th, 1989) 267

Notes

Analysis of fatty acids in equine cerebrospinal fluid using gas chromatography with electron-capture detection
by R.W. Sweeney, J. Beech, R.H. Whitlock and P.L. Castelli (Kennett Square, PA, U.S.A.) (Received May 3rd, 1989) 278

Determination of phosphoethanolamine in animal tissues by gas chromatography with flame photometric detection
by H. Kataoka, N. Sakiyama, M. Maeda and M. Makita (Okayama, Japan) (Received May 30th, 1989) 283

Rapid, quantitative method for the isolation and purification of gangliosides by LIPSEP gel chromatography
by K. Kant, A.P. Joshi and K.C. Gupta (Delhi, India) (Received May 3rd, 1989) ... 289

Simultaneous measurement of acetylcholine and dopamine releases in rat striatum under freely moving conditions with a brain dialysis method
by A. Ajima, T. Nakagawa and T. Kato (Yokohama, Japan) (Received May 10th, 1989) 297

(Continued overleaf)

Contents (continued)

Measurement of urinary free catecholamines using high-performance liquid chromatography with electrochemical detection
by B.L. Lee, K.S. Chia and C.N. Ong (Singapore, Singapore) (Received April 20th, 1989) 303

Purification of the isolated β-chain of adult human haemoglobin from its post-translational modification
by M. Castagnola, L. Cassiano, R. De Cristofaro, S. Luciani, D.V. Rossetti and R. Landolfi (Rome, Italy) (Received April 25th, 1989) 310

Determination of meprobamate in serum by alkaline hydrolysis, trimethylsilyl derivatization and detection by gas chromatography–mass spectrometry
by J. Stidman, E.H. Taylor, H.F. Simmons, J. Gandy and A.A. Pappas (Little Rock, AR, U.S.A.) (Received May 17th, 1989) 318

Characterization of the antidiarrhoeal loperamide by gas chromatography–mass spectrometry and application of the Hofmann degradation and Cope elimination reaction
by H.J. Leis (Graz, Austria) and H. Gleispach (Riyadh, Saudi Arabia) (Received May 17th, 1989) 324

Gas chromatographic and mass spectrometric analysis of articaine in urine
by S.M. Fox, S.C. Chan and D.H. Van Middlesworth (Calgary, Canada) (Received April 20th, 1989) 331

Rapid high-performance liquid chromatographic method for the determination of oxolinic acid in fish serum employing solid-phase extraction
by S.O. Hustvedt and R. Salte (Ås-NLH, Norway) and T. Benjaminsen (Oslo, Norway) (Received April 24th, 1989) 335

High-performance liquid chromatographic determination of teicoplanin in plasma: comparison with a microbiological assay
by A. Georgopoulos, M.J. Czejka, N. Starzengruber, W. Jäger and H. Lackner (Vienna, Austria) (Received April 25th, 1989) 340

Determination of velnacrine in plasma and urine by high-performance liquid chromatography
by R.S. Hsu, E.M. DiLeo, S.M. Chesson, D.B. Ellis and J.K. Wichmann (Somerville, NJ, U.S.A.) (Received May 3rd, 1989) 347

Determination of mizoribine in plasma using ion-pair high-performance liquid chromatography
by G.R. Erdmann, S.A. Gruber, M.M. McGuiggan, R.J. Cipolle and D.M. Canafax (Minneapolis, MN, U.S.A.) (Received May 1st, 1989) 354

Determination of crotamiton in plasma and urine by high-performance liquid chromatography
by A. Sioufi, N. Sandrenan and J.P. Dubois (Rueil-Malmaison, France) (Received May 10th, 1989) 361

High-performance liquid chromatographic assay for timolol in the aqueous humor of the eye
by P.-Y. Wu, M. Riegel and P.P. Ellis (Denver, CO, U.S.A.) (Received May 9th, 1989) 368

Reversed-phase high-performance liquid chromatographic assay for the determination of mezlocillin in human and rat biological samples
by G.L. Jungbluth, D.M. Janicke and W.J. Jusko (Buffalo, NY, U.S.A.) (Received May 2nd, 1989) 376

Stereoselective determination of plasma pindolol in endotoxin-pretreated rats by high-performance liquid chromatography
by R. Hasegawa (Tokyo, Japan) and M. Murai-Kushiya, T. Komuro and T. Kimura (Osaka, Japan) (Received May 16th, 1989) 381

High-performance liquid chromatography of chlorambucil prodrugs structurally related to lipids in rat plasma
by J.C. Saraiva Gonçalves, C. Razzouk, J.H. Poupaert and P. Dumont (Brussels, Belgium) (Received May 15th, 1989) 389

Determination of tacrine and its 1-hydroxy metabolite in plasma using column liquid chromatography with ultraviolet detection
by L. Ekman, B. Lindström and P. Roxin (Uppsala, Sweden) (Received May 22nd, 1989) 397

Sensitive high-performance liquid chromatographic assay for albendazole and its main metabolite albendazole sulphoxide in plasma and cerebrospinal fluid
by M. Hurtado, M.T. Medina, J. Sotelo and H. Jung (Mexico, Mexico) (Received May 22nd, 1989) 403

Column liquid chromatography of calcium channel blockers
by H. Waki and S. Ando (Tokyo, Japan) (Received May 6th, 1989) 408

High-performance liquid chromatographic method for a clozapine analogue, CGS 13429, and its N-oxide and desmethyl metabolites
by J.P. Chovan and J.D. Vermeulen (Ardsley, NY, U.S.A.) (Received May 12th, 1989) 413

Letters to the Editor

Gas chromatographic–mass spectrometric analysis of *threo*-methylphenidate enantiomers in plasma
by T. Aoyama, H. Kotaki and Y. Saitoh (Tokyo, Japan) (Received May 17th, 1989) 420

Modified high-performance liquid chromatographic assay for antipyrine and its three major metabolites in urine
by J.V. St. Peter and W.M. Awni (Minneapolis, MN, U.S.A.) (Received May 12th, 1989) 424

Related articles published in Journal of Chromatography, Vol. 478 428

Author Index 429

Subject Index 434

In articles with more than one author, the name of the author to whom correspondence should be addressed is indicated in the article heading by a 6-pointed asterisk (*)